




18

The appalling and unforgettable odor of skunks is due to a mixture of several simple thiols. Comstock Images/
Jupiterimages


醚，环氧化物，硫化物



18.1 命名和性质




Isopropyl methyl ether




Ethyl phenyl ether

If other functional groups are present, the ether part is considered an *alkoxy* substituent. For example:






p-Dimethoxybenzene



4-*tert*-Butoxy-1-cyclohexene


Like alcohols, ethers have nearly the same geometry as water. The R–O–R bonds have an approximately tetrahedral bond angle (112° in dimethyl ether), and the oxygen atom is sp^3 -hybridized.



18.2 醚的合成

- 工业制法

Diethyl ether and other simple symmetrical ethers are prepared industrially by the sulfuric acid-catalyzed reaction of alcohols. The reaction occurs by S_N2 displacement of water from a protonated ethanol molecule by the oxygen atom of a second ethanol. Unfortunately, the method is limited to use with primary alcohols because secondary and tertiary alcohols dehydrate by an E1 mechanism to yield alkenes (**Section 17.6**).




18.2 醚的合成

- 威廉森醚化反应

The Williamson Ether Synthesis


The most generally useful method of preparing ethers is by the *Williamson ether synthesis*, in which an alkoxide ion reacts with a primary alkyl halide or tosylate in an S_N2 reaction. As we saw earlier in **Section 17.2**, the alkoxide ion is normally prepared by reaction of an alcohol with a strong base such as sodium hydride, NaH.



18.2 醚的合成



- Ag_2O 作为更温和的碱使用

A useful variation of the Williamson synthesis involves silver oxide, Ag_2O , as a mild base rather than NaH . Under these conditions, the free alcohol reacts directly with alkyl halide, so there is no need to preform the metal alkoxide intermediate. Sugars react particularly well; glucose, for example, reacts with excess iodomethane in the presence of Ag_2O to generate a pentaether in 85% yield.



18.2 醚的合成

- 注意选择合适的醇和卤化物



Problem 18.3

How would you prepare the following ethers using a Williamson synthesis?



- (a) Methyl propyl ether (b) Anisole (methyl phenyl ether)
(c) Benzyl isopropyl ether (d) Ethyl 2,2-dimethylpropyl ether





18.2 醚的合成

Alkoxymercuration of Alkenes


We saw in **Section 8.4** that alkenes react with water in the presence of mercuric acetate to yield a hydroxymercuration product. Subsequent treatment with NaBH_4 breaks the C–Hg bond and yields the alcohol. A similar **alkoxymercuration** reaction occurs when an alkene is treated with an *alcohol* in the presence of mercuric acetate or, even better, mercuric trifluoroacetate, $(\text{CF}_3\text{CO}_2)_2\text{Hg}$. Demercuration by reaction with NaBH_4 then yields an ether. The net result is Markovnikov addition of the alcohol to the alkene.



18.3 醚的反应：酸催化切断




Ethers with a tertiary, benzylic, or allylic group cleave by either an $\text{S}_{\text{N}}1$ or $\text{E}1$ mechanism because these substrates can produce stable intermediate carbocations. These reactions are often fast and take place at moderate temperatures. *tert*-Butyl ethers, for example, react by an $\text{E}1$ mechanism on treatment with trifluoroacetic acid at 0°C . We'll see in **Section 26.7** that the reaction is often used in the laboratory synthesis of peptides.



18.3 醚的反应：酸催化切断

Predicting the Product of an Ether Cleavage Reaction


Predict the products of the following reaction:





Strategy

Identify the substitution pattern of the two groups attached to oxygen—in this case a tertiary alkyl group and a primary alkyl group. Then recall the guidelines for ether cleavages. An ether with only primary and secondary alkyl groups usually undergoes cleavage by $\text{S}_{\text{N}}2$ attack of a nucleophile on the less hindered alkyl group, but an ether with a tertiary alkyl group usually undergoes cleavage by an $\text{S}_{\text{N}}1$ mechanism. In this case, an $\text{S}_{\text{N}}1$ cleavage of the tertiary C–O bond will occur, giving 1-propanol and a tertiary alkyl bromide. In addition, a competitive $\text{E}1$ reaction leading to alkene might occur.


Solution



18.4 克莱森重排




A similar rearrangement takes place with allyl vinyl ethers, leading to a so-called γ,δ -unsaturated ketone or aldehyde.




18.4 克莱森重排

Like the Diels–Alder reaction discussed in **Sections 14.4 and 14.5**, the Claisen rearrangement reaction takes place in a single step through a **pericyclic mechanism** in which a reorganization of bonding electrons occurs through a six-membered, cyclic transition state. The 6-allyl-2,4-cyclohexadienone intermediate then isomerizes to *o*-allylphenol (**Figure 18.1**).





Problem 18.10


What product would you expect from Claisen rearrangement of 2-butenyl phenyl ether?




18.5 环醚



In the laboratory, as we saw in **Section 8.7**, epoxides are usually prepared by treatment of an alkene with a peroxyacid (RCO_3H), typically *m*-chloroperoxybenzoic acid.



18.5 环醚





Problem 18.11

Reaction of *cis*-2-butene with *m*-chloroperoxybenzoic acid yields an epoxide different from that obtained by reaction of the *trans* isomer. Explain.





18.6 环醚的开环反应

- 酸催化开环




18.6 环醚的开环反应

- 酸催化开环：不对称底物通常给出混合产物




18.6 环醚的开环反应

- 碱性条件开环




Base-catalyzed epoxide opening is a typical S_N2 reaction in which attack of the nucleophile takes place at the less hindered epoxide carbon. For example, 1,2-epoxypropane reacts with ethoxide ion exclusively at the less highly substituted, primary, carbon to give 1-ethoxy-2-propanol.





18.6 环醚的开环反应

- 碱性条件开环




Problem 18.14

Predict the major product of the following reactions:



18.7 冠醚



18-Crown-6 ether

The importance of crown ethers derives from their ability to sequester specific metal cations in the center of the polyether cavity. 18-Crown-6, for example, binds strongly with potassium ion. As a result, a solution of 18-crown-6 in a nonpolar organic solvent will dissolve many potassium salts. Potassium permanganate, KMnO_4 , dissolves in toluene in the presence of 18-crown-6, for instance, and the resulting solution is a valuable reagent for oxidizing alkenes.



18.7 冠醚

- 1987 年诺贝尔化学奖：主客体化学

The Nobel Prize in Chemistry 1987




Photo from the Nobel Foundation archive.

Donald J. Cram

Prize share: 1/3



Photo from the Nobel Foundation archive.

Jean-Marie Lehn

Prize share: 1/3




Photo from the Nobel Foundation archive.


Charles J. Pedersen

Prize share: 1/3


The Nobel Prize in Chemistry 1987 was awarded jointly to Donald J. Cram, Jean-Marie Lehn and Charles J. Pedersen "for their development and use of molecules with structure-specific interactions of high selectivity."




18.8 硫醇和硫醚





Ethanethiol






Cyclohexanethiol



m-Mercaptobenzoic acid



18.8 硫醇和硫醚




18.8 硫醇和硫醚

- 课堂练习

Problem 18.17

2-Butene-1-thiol is one component of skunk spray. How would you synthesize this substance from methyl 2-butenate? From 1,3-butadiene?



18.9 波谱分析





Figure 18.5 The ^1H NMR spectrum of 1,2-epoxypropane.

Ether carbon atoms also exhibit a downfield shift in the ^{13}C NMR spectrum, where they usually absorb in the 50 to 80 δ range. For example, the carbon atoms next to oxygen in methyl propyl ether absorb at 58.5 and 74.8 δ . Similarly, the methyl carbon in anisole absorbs at 54.8 δ .




18.9 波谱分析

- 课堂练习

Problem 18.18

The ^1H NMR spectrum shown is that of a cyclic ether with the formula $\text{C}_4\text{H}_8\text{O}$. Propose a structure.



作业

- 18.47, 18.53, 18.55, 18.58, 18.61

